

# UNIT GROUPS OF CUBE RADICAL ZERO COMMUTATIVE COMPLETELY PRIMARY FINITE RINGS

CHITENG'A JOHN CHIKUNJI

Received 1 July 2004

A completely primary finite ring is a ring  $R$  with identity  $1 \neq 0$  whose subset of all its zero divisors forms the unique maximal ideal  $J$ . Let  $R$  be a commutative completely primary finite ring with the unique maximal ideal  $J$  such that  $J^3 = (0)$  and  $J^2 \neq (0)$ . Then  $R/J \cong GF(p^r)$  and the characteristic of  $R$  is  $p^k$ , where  $1 \leq k \leq 3$ , for some prime  $p$  and positive integer  $r$ . Let  $R_o = GR(p^{kr}, p^k)$  be a Galois subring of  $R$  and let the annihilator of  $J$  be  $J^2$  so that  $R = R_o \oplus U \oplus V$ , where  $U$  and  $V$  are finitely generated  $R_o$ -modules. Let nonnegative integers  $s$  and  $t$  be numbers of elements in the generating sets for  $U$  and  $V$ , respectively. When  $s = 2$ ,  $t = 1$ , and the characteristic of  $R$  is  $p$ ; and when  $t = s(s+1)/2$ , for any fixed  $s$ , the structure of the group of units  $R^*$  of the ring  $R$  and its generators are determined; these depend on the structural matrices  $(a_{ij})$  and on the parameters  $p, k, r$ , and  $s$ .

## Notations

Throughout this paper,  $R$  will denote a finite ring, unless otherwise stated,  $J$  will denote the Jacobson radical of  $R$ , and we will denote the Galois ring  $GR(p^{nr}, p^n)$  of characteristic  $p^n$  and order  $p^{nr}$  by  $R_o$ , for some prime  $p$ , and positive integers  $n, r$ .

We denote the group of units of  $R$  by  $R^*$  and a cyclic group of order  $\pi$  by  $\epsilon(\pi)$ . If  $g$  is an element of  $R^*$ , then  $o(g)$  denotes its order, and  $\langle g \rangle$  denotes the cyclic group generated by  $g$ . Furthermore, for a subset  $A$  of  $R$  or  $R^*$ ,  $|A|$  will denote the number of elements in  $A$ . The ring of integers modulo the number  $n$  will be denoted by  $\mathbb{Z}_n$ , and the characteristic of  $R$  will be denoted by  $\text{char } R$ .

## 1. Introduction

In [6], Fuchs asked for a characterization of abelian groups which could be groups of units of a ring. This question was noted to be too general for a complete answer [12], and a natural course is to restrict the classes of groups or rings to be considered.

Let  $R$  be a ring and let  $R^*$  denote its multiplicative group of unit elements. All local rings  $R$  with  $R^*$  cyclic were determined by Gilmer [8] and this case was also considered by Ayoub [1] (also proofs are given in [10, 11]). Pearson and Schneider have found all

$R$  where  $R^*$  is generated by two elements. Clark [4] has investigated  $R^*$  where the ideals form a chain and has shown that if  $p \geq 3$ ,  $n \geq 2$ , and  $r \geq 2$ , then the units of the Galois ring  $GR(p^{nr}, p^n)$  are a direct sum of a cyclic group of order  $p^r - 1$  and  $r$  cyclic groups of order  $p^n - 1$  (this was also done independently by Raghavendran [11]). In fact, Raghavendran described the structure of the multiplicative group of every Galois ring. Stewart in [12] considered a related problem to that asked by Fuchs [6] by proving that for a given finite group  $G$  (not necessarily abelian), there are, up to isomorphism, only finitely many directly indecomposable finite rings having group of units isomorphic to  $G$ .

Ganske and McDonald [7] provided a solution for  $R^*$  when the local ring  $R$  has Jacobson radical  $J$  such that  $J^2 = (0)$  by showing that

$$R^* = \left( \bigoplus_{i=1}^{nt} \epsilon(p) \right) \oplus \epsilon(|K| - 1), \quad (1.1)$$

where  $n = \dim_K(J/J^2)$ ,  $|K| = p^t$ , and  $\epsilon(\pi)$  denotes the cyclic group of order  $\pi$ .

In [5], Dolzan found all nonisomorphic rings with a group of units isomorphic to a group  $G$  with  $n$  elements, where  $n$  is a power of a prime or any product of prime powers, not divisible by 4; and also found all groups with  $n$  elements which can be groups of units of a finite ring, a contribution to Stewart's problem [12]. More recently, X.-D. Hou et al. gave an algorithmic method for computing the structure of the group of units of a finite commutative chain ring and further strengthening the known result by listing a set of linearly independent generators for the group of units.

The present paper focuses on the group of units  $R^*$  of a commutative completely primary finite ring  $R$  with unique maximal ideal  $J$  such that  $R/J \cong GF(p^r)$ ,  $J^3 = (0)$ , and  $J^2 \neq (0)$  so that the characteristic of  $R$  is  $p^k$ , where  $1 \leq k \leq 3$ ; and further identifies sets of generators for  $R^*$ .

In particular, let  $R_o = GR(p^{kr}, p^k)$  be a Galois subring of  $R$  and let the annihilator of  $J$  be  $J^2$  so that  $R = R_o \oplus U \oplus V$ , where  $U$  and  $V$  are finitely generated  $R_o$ -modules. Let nonnegative integers  $s$  and  $t$  be numbers of elements in the generating sets for  $U$  and  $V$ , respectively. When  $s = 2$ ,  $t = 1$ , and  $\text{char } R = p$ , and when  $t = s(s+1)/2$ , for any fixed  $s$ , the structure of the group of units  $R^*$  of the ring  $R$  and its generators have been determined; these depend on the structural matrices  $(a_{ij})$  and on the parameters  $p$ ,  $k$ ,  $r$ , and  $s$ .

## 2. Preliminaries

We refer the reader to [2] for the general background of completely primary finite rings  $R$  with maximal ideals  $J$  such that  $J^3 = \{0\}$  and  $J^2 \neq \{0\}$ . Let  $R$  be a completely primary finite ring with maximal ideal  $J$  such that  $J^3 = (0)$  and  $J^2 \neq (0)$ . Then  $R$  is of order  $p^{nr}$  and the residue field  $R/J$  is a finite field  $GF(p^r)$ , for some prime  $p$  and positive integers  $n$ ,  $r$ . The characteristic of  $R$  is  $p^k$ , where  $k$  is an integer such that  $1 \leq k \leq 3$ . Let  $GR(p^{kr}, p^k)$  be the Galois ring of characteristic  $p^k$  and order  $p^{kr}$ , that is,  $GR(p^{kr}, p^k) = \mathbb{Z}_{p^k}[x]/(f)$ , where  $f \in \mathbb{Z}_{p^k}[x]$  is a monic polynomial of degree  $r$  whose image in  $\mathbb{Z}_p[x]$  is irreducible. Then, it can be deduced from the main theorem in [4] that  $R$  has a coefficient subring  $R_o$  of the form  $GR(p^{kr}, p^k)$  which is clearly a maximal Galois subring of  $R$ . Moreover, there

exist elements  $m_1, m_2, \dots, m_h \in J$  and automorphisms  $\sigma_1, \dots, \sigma_h \in \text{Aut}(R_o)$  such that

$$R = R_o \oplus \sum_{i=1}^h R_o m_i \tag{2.1}$$

(as  $R_o$ -modules),  $m_i r = r^{\sigma_i} m_i$ , for every  $r \in R_o$  and any  $i = 1, \dots, h$ . Further,  $\sigma_1, \dots, \sigma_h$  are uniquely determined by  $R$  and  $R_o$ . The maximal ideal of  $R$  is

$$J = pR_o \oplus \sum_{i=1}^h R_o m_i. \tag{2.2}$$

It is worth noting that  $R$  contains an element  $b$  of multiplicative order  $p^r - 1$  and that  $R_o = \mathbb{Z}_{p^k}[b]$  (see, e.g., [2, Result 1.3]).

The following results will be useful.

**PROPOSITION 2.1.** *Let  $R$  be a completely primary finite ring (not necessarily commutative). Then the group of units  $R^*$  of  $R$  contains a cyclic subgroup  $\langle b \rangle$  of order  $p^r - 1$ , and  $R^*$  is a semidirect product of  $1 + J$  and  $\langle b \rangle$ .*

*Proof.* Obviously, the group of units  $R^*$  of  $R$  is  $R - J$ ,  $|R^*| = p^{(n-1)r}(p^r - 1)$ , and  $\phi : R \rightarrow R/J$  induces a surjective multiplicative group homomorphism  $\varphi : R^* \rightarrow (R/J)^*$ . Since  $\ker \phi = J$ , we have  $\ker \varphi = 1 + J$ . In particular,  $1 + J$  is a normal subgroup of  $R^*$ .

Let  $\langle \beta \rangle = (R/J)^*$ , and let  $b_o \in \varphi^{-1}(\beta)$ . Then, the multiplicative order of  $b_o$  is a multiple of  $p^r - 1$  and a divisor of  $|R - J| = p^{nr} - p^{(n-1)r} = p^{(n-1)r}(p^r - 1)$ ; hence, of the form  $p^s(p^r - 1)$ . But then  $b = b_o^{p^s}$  has multiplicative order  $p^r - 1$  and  $\varphi(b_o^{p^s}) = \beta^{p^s}$ , which is still a generator of  $(R/J)^*$ , since  $(p^s, p^r - 1) = 1$ .

Finally, since  $|R^*| = |1 + J| \cdot |\langle b \rangle|$ , and  $(1 + J) \cap \langle b \rangle = 1$ , we have  $R^* = (1 + J) \cdot \langle b \rangle$ , hence,  $R^* = (1 + J) \times_{\theta} \langle b \rangle$ , a semidirect product. □

**PROPOSITION 2.2.** *Let  $R$  be a completely primary finite ring (not necessarily commutative). Then the group of units  $R^*$  is solvable.*

*Proof.* That  $R^*$  is a solvable group follows from the fact that  $1 + J$  is a normal  $p$ -subgroup of  $R^*$ , and  $R^*/(1 + J)$  is cyclic. □

**LEMMA 2.3.** *Let  $R$  be a completely primary finite ring (not necessarily commutative). If  $G$  is a subgroup of  $R^*$  of order  $p^r - 1$ , then  $G$  is conjugate to  $\langle b \rangle$  in  $R^*$ .*

*Proof.* This follows from key properties of  $p$ -solvable groups contained in the variation of Sylow's theorem, due to Philip Hall, since the order of  $G$  is prime to its index in  $R^*$  (see, e.g., [9, Theorem 8.2 page 25]). □

**PROPOSITION 2.4.** *Let  $R$  be a completely primary finite ring (not necessarily commutative). If  $R^*$  contains a normal subgroup of order  $p^r - 1$ , then the set  $K_o = \langle b \rangle \cup \{0\}$  is contained in the center of the ring  $R$ .*

*Proof.* By Lemma 2.3,  $\langle b \rangle$  is normal in  $R^*$  and since  $1 + J$  is a normal subgroup of  $R^*$  with  $|\langle b \rangle \cap (1 + J)| = 1$ , it follows that  $\langle b \rangle$  and  $1 + J$  commute elementwise. Hence,  $b$  lies in the center of  $R$ . □

PROPOSITION 2.5. *Let  $R$  be a completely primary finite ring. Then,  $(1 + J^i)/(1 + J^{i+1}) \cong J^i/J^{i+1}$  (the left-hand side as a multiplicative group and the right-hand side as an additive group).*

*Proof.* Consider the map

$$\eta : (1 + J^i)/(1 + J^{i+1}) \longrightarrow J^i/J^{i+1} \tag{2.3}$$

defined by

$$(1 + x)(1 + J^{i+1}) \longrightarrow x + J^{i+1}. \tag{2.4}$$

Then it is easy to see that  $\eta$  is an isomorphism. □

*Remark 2.6* (see [3, Result 2.7]). Let  $R$  be a completely primary finite ring of characteristic  $p^k$  and with Jacobson radical  $J$ . Let  $R_o$  be a Galois subring of  $R$ . If  $m \in J$  and  $p^t$  is the additive order of  $m$ , for some positive integer  $t$ , then  $|R_o m| = p^{tr}$ .

*Proof.* Apply the fact that

$$R_o m \cong R_o/p^t R_o. \tag{2.5}$$

□

Now let  $R$  be a commutative completely primary finite ring with maximal ideal  $J$  such that  $J^3 = (0)$  and  $J^2 \neq (0)$ . In [2], the author gave constructions describing these rings for each characteristic and for details, we refer the reader to [2, Sections 4 and 6].

If  $R$  is a commutative completely primary finite ring with maximal ideal  $J$  such that  $J^3 = (0)$  and  $J^2 \neq (0)$ , then from Constructions A and B [2],

$$R = R_o \oplus U \oplus V \oplus W, \tag{2.6}$$

$$J = pR_o \oplus U \oplus V \oplus W, \tag{2.7}$$

where the  $R_o$ -modules  $U$ ,  $V$ , and  $W$  are finitely generated. The structure of  $R$  is characterized by the invariants  $p, n, r, d, s, t$ , and  $\lambda$ ; and the linearly independent matrices  $(a_{ij}^k)$  defined in the multiplication. Let  $\text{ann}(J)$  denote the two-sided annihilator of  $J$  in  $R$ . Notice that since  $J^2 \subseteq \text{ann}(J)$ , we can write  $R = R_o \oplus U \oplus M$ , and hence,  $J = pR_o \oplus U \oplus M$ , where  $M = V \oplus W$ , and the multiplication in  $R$  may be written accordingly. It is therefore easy to see that the description of rings of this type reduces to the case where  $\text{ann}(J)$  coincides with  $J^2$ . Therefore, when investigating the structure of the group of units of this type of rings for a given order, say  $p^{nr}$ , where  $\text{ann}(J)$  does not coincide with  $J^2$ , we will first write all the rings of this type of order  $\leq p^{nr}$ , where  $\text{ann}(J)$  coincides with  $J^2$ .

In what follows, we assume that  $\text{ann}(J) = J^2$ .

Let  $R_o = GR(p^{kr}, p^k) (1 \leq k \leq 3)$  and let nonnegative integers  $s$  and  $t$  be numbers of elements in the generating sets  $\{u_1, \dots, u_s\}$  and  $\{v_1, \dots, v_t\}$  for finitely generated  $R_o$ -modules  $U$  and  $V$ , respectively, where  $t \leq s(s + 1)/2$ . Assume that  $u_1, u_2, \dots, u_s$  and  $v_1, \dots, v_t$  are commuting indeterminates. Then  $R = R_o \oplus U \oplus V$ .

By Proposition 2.1, and since  $R$  is commutative,

$$R^* = \langle b \rangle \cdot (1 + J) \cong \langle b \rangle \times (1 + J), \tag{2.8}$$

a direct product.

Again, notice that since  $R$  is of order  $p^{nr}$  and  $R^* = R - J$ , it is easy to see that  $|R^*| = p^{(n-1)r}(p^r - 1)$  and  $|1 + J| = p^{(n-1)r}$ , so that  $1 + J$  is an abelian  $p$ -group. Thus,  $R^* \cong (\text{abelian } p\text{-group}) \times (\text{cyclic group of order } |R/J| - 1)$ .

Our goal is to determine the structure and identify a set of generators of the multiplicative abelian  $p$ -group  $1 + J$ .

### 3. The group $1 + J$

Now let  $R$  be a commutative completely primary finite ring with maximal ideal  $J$  such that  $J^3 = (0)$  and  $J^2 \neq (0)$ . Let  $1 + J$  be the abelian  $p$ -subgroup of the unit group  $R^*$ .

The group  $1 + J$  has a filtration  $1 + J \supset 1 + J^2 \supset 1 + J^3 = \{1\}$  with filtration quotients  $(1 + J)/(1 + J^2)$  and  $(1 + J^2)/\{1\} = 1 + J^2$  isomorphic to the additive groups  $J/J^2$  and  $J^2$ , respectively.

*Remark 3.1.* Notice that  $1 + J^2$  is a normal subgroup of  $1 + J$ . But, in general,  $1 + J$  does not have a subgroup which is isomorphic to the quotient  $(1 + J)/(1 + J^2)$  as may be illustrated by the following example.

*Example 3.2.* Let  $R = \mathbb{Z}_{p^3}$ , where  $p$  is an odd prime. Then  $J = p\mathbb{Z}_{p^3}$ ,  $\text{ann}(J) = J^2$ , and  $1 + J \cong \mathbb{Z}_{p^2}$ ,  $1 + J^2 \cong \mathbb{Z}_p$ ,  $(1 + J)/(1 + J^2) \cong \mathbb{Z}_p$ .

*Remark 3.3.* In view of the above remark and example, we investigate the structure of  $1 + J$  by considering various subgroups of  $1 + J$ .

**3.1. The case when  $s = 2, t = 1$ , and  $\text{char } R = p$ .** Suppose  $s = 2, t = 1$ , and  $\text{char } R = p$ . Let  $R_o = \mathbb{F}_q = GF(p^r)$ , the Galois field of  $q = p^r$  elements. Then

$$R = \mathbb{F}_q \oplus \mathbb{F}_q u_1 \oplus \mathbb{F}_q u_2 \oplus \mathbb{F}_q v, \tag{3.1}$$

the Jacobson radical

$$J = \mathbb{F}_q u_1 \oplus \mathbb{F}_q u_2 \oplus \mathbb{F}_q v, \tag{3.2}$$

$$J^2 = \mathbb{F}_q v. \tag{3.3}$$

The multiplication in  $R$  is given by

$$u_1^2 = a_{11}v, \quad u_1 u_2 = u_2 u_1 = a_{12}v, \quad u_2^2 = a_{22}v, \tag{3.4}$$

where  $a_{ij} \in \mathbb{F}_q$ . The elements  $a_{ij}$  form a nonzero symmetric matrix

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \tag{3.5}$$

since  $J^2 \neq (0)$ .

Since  $R^*$  is a direct product of the cyclic group  $\langle b \rangle$  of order  $p^r - 1$  and the group  $1 + J$  of order  $p^{3r}$ , it suffices to determine the structure of  $1 + J$ .

In this case,

$$1 + J = 1 + \mathbb{F}_q u_1 \oplus \mathbb{F}_q u_2 \oplus \mathbb{F}_q v, \tag{3.6}$$

and since  $s$  and  $t$  are fixed, the structure of  $1 + J$  now depends on the prime  $p$ , the integer  $r$ , and the structural matrix  $\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ . We investigate this by considering cases depending on the type of the structural matrix.

Let  $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_r$  be elements of  $\mathbb{F}_q$  with  $\varepsilon_1 = 1$  so that  $\overline{\varepsilon_1}, \overline{\varepsilon_2}, \dots, \overline{\varepsilon_r}$  form a basis for  $\mathbb{F}_q$  regarded as a vector space over its prime subfield  $\mathbb{F}_p$ .

*Case (i).* Suppose that  $\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix}$ , with  $a \neq 0$ . Then

$$1 + J \cong \begin{cases} \mathbb{Z}_4^r \times \mathbb{Z}_2^r, & \text{if } \text{char } R = 2, \\ \mathbb{Z}_p^r \times \mathbb{Z}_p^r \times \mathbb{Z}_p^r, & \text{if } \text{char } R = p \neq 2. \end{cases} \tag{3.7}$$

To see this, we consider the two cases separately. So, suppose that  $p = 2$ . We first note the following results:

$$1 + \varepsilon_i u_1 \in 1 + J, \quad (1 + \varepsilon_i u_1)^4 = 1, \quad (1 + \varepsilon_i u_2)^2 = 1, \quad g^4 = 1, \quad \forall g \in 1 + J. \tag{3.8}$$

For positive integers  $k_i, l_i$ , with  $k_i \leq 4, l_i \leq 2$ , we assert that

$$\prod_{i=1}^r \left\{ (1 + \varepsilon_i u_1)^{k_i} \right\} \cdot \prod_{i=1}^r \left\{ (1 + \varepsilon_i u_2)^{l_i} \right\} = 1 \tag{3.9}$$

will imply  $k_i = 4$  for all  $i = 1, \dots, r$ ; and  $l_i = 2$  for all  $i = 1, \dots, r$ .

If we set  $F_i = \{(1 + \varepsilon_i u_1)^k \mid k = 1, \dots, 4\}$  for all  $i = 1, \dots, r$ ; and  $G_i = \{(1 + \varepsilon_i u_2)^l \mid l = 1, 2\}$  for all  $i = 1, \dots, r$ , we see that  $F_i, G_i$  are all cyclic subgroups of the group  $1 + J$  and that these are of the precise orders indicated by their definition. The argument above will show that the product of  $2r$  subgroups  $F_i$  and  $G_i$  is direct. So, their product will exhaust the group  $1 + J$ .

When  $p$  is an odd prime, we have to consider the equation

$$\prod_{i=1}^r \left\{ (1 + \varepsilon_i u_1)^{k_i} \right\} \cdot \prod_{i=1}^r \left\{ (1 + \varepsilon_i u_2)^{l_i} \right\} \cdot \prod_{i=1}^r \left\{ (1 + \varepsilon_i v)^{m_i} \right\} = 1 \tag{3.10}$$

and as each element in  $1 + J$  raised to the power  $p$  equals 1, we see that  $1 + J$  will be an elementary abelian group.

*Case (ii).* Suppose that  $\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} 0 & a \\ a & 0 \end{pmatrix}$ , with  $a \neq 0$ . Then

$$1 + J \cong \mathbb{Z}_p^r \times \mathbb{Z}_p^r \times \mathbb{Z}_p^r, \tag{3.11}$$

for every  $p = \text{char } R$ . In this case, we consider the equation

$$\prod_{i=1}^r \{(1 + \varepsilon_i u_1)^{k_i}\} \cdot \prod_{i=1}^r \{(1 + \varepsilon_i u_2)^{l_i}\} \cdot \prod_{i=1}^r \{(1 + \varepsilon_i v)^{m_i}\} = 1 \tag{3.12}$$

and the integers  $k_i, l_i, m_i$  will imply  $k_i = l_i = m_i = p$  for all  $i = 1, \dots, r$ .

If we set  $F_i = \{(1 + \varepsilon_i u_1)^k | k = 1, \dots, p\}$  for all  $i = 1, \dots, r$ ;  $G_i = \{(1 + \varepsilon_i u_2)^l | l = 1, \dots, p\}$  for all  $i = 1, \dots, r$ ; and  $H_i = \{(1 + \varepsilon_i v)^m | m = 1, \dots, p\}$  for all  $i = 1, \dots, r$ , we see that  $F_i, G_i$ , and  $H_i$  are all cyclic subgroups of the group  $1 + J$  and that these are all of order  $p$ . The product of the  $3r$  subgroups  $F_i, G_i$ , and  $H_i$  is direct. So, their product will exhaust the group  $1 + J$ .

Case (iii). Suppose now that  $\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} a & b \\ b & 0 \end{pmatrix}$ , with  $a$  and  $b$  being nonzero. Then

$$1 + J \cong \begin{cases} \mathbb{Z}_4^r \times \mathbb{Z}_2^r, & \text{if } \text{char } R = 2, \\ \mathbb{Z}_p^r \times \mathbb{Z}_p^r \times \mathbb{Z}_p^r, & \text{if } \text{char } R = p \neq 2. \end{cases} \tag{3.13}$$

The argument is similar to that in Case (i).

Case (iv). Suppose  $\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$ , with  $a$  and  $b$  being nonzero. Then  $u_1^2 = av, u_2^2 = bv$ , and  $u_1 u_2 = u_2 u_1 = 0$ .

If  $\text{char } R = p \neq 2$ , then  $o(1 + \varepsilon_i u_1) = o(1 + \varepsilon_i u_2) = p (i = 1, \dots, r)$ . Moreover, for every  $i = 1, \dots, r$ ,  $\langle 1 + \varepsilon_i u_1 \rangle \cap \langle 1 + \varepsilon_i u_2 \rangle = \{1\}$ . Also,  $o(1 + \varepsilon_i v) = p$ , and the element  $1 + \varepsilon_i v (i = 1, \dots, r)$  generates a cyclic subgroup of order  $p$ .

If  $\text{char } R = 2$ , then in  $1 + J$ , we see that  $o(1 + \varepsilon_i u_1) = 4$  and for each  $\varepsilon_i$ , by considering the element  $1 + \varepsilon_i u_1 + \varepsilon_i u_2 + \varepsilon_i v$  of order 2, one obtains the direct product

$$1 + J = \prod_{i=1}^r \langle 1 + \varepsilon_i u_1 \rangle \times \prod_{i=1}^r \langle 1 + \varepsilon_i u_1 + \varepsilon_i u_2 + \varepsilon_i v \rangle. \tag{3.14}$$

Hence,

$$1 + J \cong \begin{cases} \mathbb{Z}_4^r \times \mathbb{Z}_2^r, & \text{if } \text{char } R = 2, \\ \mathbb{Z}_p^r \times \mathbb{Z}_p^r \times \mathbb{Z}_p^r, & \text{if } \text{char } R = p \neq 2. \end{cases} \tag{3.15}$$

Case (v). Finally, suppose that  $\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} a & b \\ b & c \end{pmatrix}$ , with  $a, b$ , and  $c$  being nonzero. Then  $u_1^2 = av, u_2^2 = cv$ , and  $u_1 u_2 = u_2 u_1 = bv$ . In this case, it is easy to verify that

$$1 + J \cong \begin{cases} \mathbb{Z}_4^r \times \mathbb{Z}_2^r, & \text{if } \text{char } R = 2, \\ \mathbb{Z}_p^r \times \mathbb{Z}_p^r \times \mathbb{Z}_p^r, & \text{if } \text{char } R = p \neq 2. \end{cases} \tag{3.16}$$

The number of cases involved in determining the structure of  $1 + J$  for larger values of  $s$  and for  $t < s(s + 1)/2$  compels us to investigate the problem by considering the extreme case when the invariant  $t = s(s + 1)/2$ , and to leave the other cases for subsequent work.

**3.2. The case when  $t = s(s + 1)/2$ , for  $s$  fixed.** Suppose that  $t = s(s + 1)/2$  for a fixed non-negative integer  $s$ . Let  $u_1, u_2, \dots, u_s$  be commuting indeterminates over the Galois ring  $R_o = GR(p^{kr}, p^k)$ , where  $1 \leq k \leq 3$ . Then it is easy to verify that

$$R = R_o \oplus \sum_{i=1}^s R_o u_i \oplus \sum_{i,j=1}^s R_o u_i u_j, \tag{3.17}$$

where

$$u_i u_j = u_j u_i, \quad u_i^3 = u_i^2 u_j = u_i u_j^2 = 0, \quad \text{for every } i, j = 1, \dots, s, \tag{3.18}$$

is a commutative completely primary finite ring with Jacobson radical

$$J = pR_o \oplus \sum_{i=1}^s R_o u_i \oplus \sum_{i,j=1}^s R_o u_i u_j; \tag{3.19}$$

$$J^2 = pR_o \oplus \sum_{i,j=1}^s R_o u_i u_j \quad \text{or} \quad J^2 = p^2 R_o \oplus \sum_{i,j=1}^s R_o u_i u_j; \quad J^3 = (0). \tag{3.20}$$

In this case, the linearly independent matrices  $(a_{ij}^k)$  defined in the multiplication of  $R$  are the  $t = s(s + 1)/2, s \times s$  symmetric matrices with 1's in the  $(i, j)$ th and  $(j, i)$ th positions, and zeros elsewhere.

It follows clearly that

$$1 + J = 1 + pR_o \oplus \sum_{i=1}^s R_o u_i \oplus \sum_{i,j=1}^s R_o u_i u_j, \tag{3.21}$$

and it can easily be deduced that every element  $x$  of  $1 + J$  has a unique expression of the form

$$x = 1 + p a_o + \sum_{i=1}^s a_i u_i + \sum_{i,j=1}^s a_{ij} u_i u_j, \tag{3.22}$$

where  $a_o, a_i, a_{ij} = a_{ji}$  are in  $K = R_o/pR_o$ .

Let  $s$  be a fixed nonnegative integer and suppose that  $t = s(s + 1)/2$ . If  $\text{char } R = p$ , then

$$|R| = p^{((s^2+3s+2)/2)r}, \quad |J| = p^{((s^2+3s)/2)r} \tag{3.23}$$

because  $|R_o u_i| = p^r$  (for each  $i = 1, \dots, s$ ) and  $|R_o u_i u_j| = p^r$  (for  $i, j = 1, \dots, s$ ); thus

$$|1 + J| = p^{((s^2+3s)/2)r}. \tag{3.24}$$

If  $\text{char } R = p^2$ , then

$$|R| = p^{((s^2+5s+4)/2)r}, \quad |J| = p^{((s^2+5s+2)/2)r} \tag{3.25}$$



because  $|R_o| = p^{2r}$ ,  $|pR_o| = p^r$ ,  $|R_o u_i| = p^{2r}$ , if  $pu_i \neq 0$  (for each  $i = 1, \dots, s$ ) and  $|R_o u_i u_j| = p^r$  (for  $i, j = 1, \dots, s$ ) (see Remark 2.6), and thus

$$|1 + J| = p^{((s^2+5s+2)/2)r}. \tag{3.26}$$

Finally, if  $\text{char } R = p^3$ , then

$$|R| = p^{((s^2+5s+6)/2)r}, \quad |J| = p^{((s^2+5s+4)/2)r} \tag{3.27}$$

because  $|R_o| = p^{3r}$ ,  $|pR_o| = p^{2r}$  and if  $pu_i \neq 0$ ,  $|R_o u_i| = p^{2r}$  (because  $p^2 u_i = 0$ ) (for each  $i = 1, \dots, s$ ) and  $|R_o u_i u_j| = p^r$  (for  $i, j = 1, \dots, s$ ) (see Remark 2.6 and also because  $pu_i u_j = 0$ ), and hence,

$$|1 + J| = p^{((s^2+5s+4)/2)r}. \tag{3.28}$$

**PROPOSITION 3.4.** *If  $\text{char } R = p^k$ , where  $k = 2$  or  $3$ , then  $1 + J$  contains  $1 + pR_o$  as its subgroup.*

*Proof.* We only show the case for  $\text{char } R = p^2$ , the other case follows easily from this. Now, each element of  $1 + pR_o$  is of the form  $1 + pr$ , for every  $r \in R_o$ , and for any two elements  $1 + pr_1$  and  $1 + pr_2$ , we have

$$(1 + pr_1)(1 + pr_2) = 1 + p(r_1 + r_2) \tag{3.29}$$

which is clearly an element of  $1 + pR_o$ . □

**PROPOSITION 3.5.** *For each pair  $u_i, u_j$  with  $i \neq j$  and  $u_i u_j = u_j u_i$ ,  $1 + R_o u_i u_j$  is a subgroup of  $1 + J$ .*

*Proof.* It is easy to see that  $1 + R_o u_i u_j$  is a subgroup of  $1 + J$  because for any two elements  $1 + r_1 u_i u_j$  and  $1 + r_2 u_i u_j$  in  $1 + R_o u_i u_j$ , we have

$$(1 + r_1 u_i u_j)(1 + r_2 u_i u_j) = 1 + (r_1 + r_2)u_i u_j \in 1 + R_o u_i u_j \tag{3.30}$$

since  $(u_i u_j)^2 = 0$ . □

**PROPOSITION 3.6.** *For every  $i = 1, \dots, s$ ,  $1 + R_o u_i + R_o u_i^2$  is a subgroup of  $1 + J$ .*

*Proof.* Obviously,

$$(1 + r_1 u_i + r_2 u_i^2)(1 + s_2 u_i + s_2 u_i^2) = 1 + (r_1 + s_1)u_i + (r_1 s_1 + r_2 + s_2)u_i^2 \tag{3.31}$$

lies in  $1 + R_o u_i + R_o u_i^2$ , for any pair  $1 + r_1 u_i + r_2 u_i^2$  and  $1 + s_2 u_i + s_2 u_i^2$  of elements in  $1 + R_o u_i + R_o u_i^2$ . □

In view of Remark 2.6 and Propositions 3.4, 3.5, and 3.6, we may now state the following.

PROPOSITION 3.7. *Let  $1 + pR_o$ ,  $1 + R_o u_i + R_o u_i^2$ , and  $1 + R_o u_i u_j$  be the subgroups of  $1 + J$  defined above. Then*

$$|1 + pR_o| = \begin{cases} p^r, & \text{if char } R = p^2, \\ p^{2r}, & \text{if char } R = p^3, \end{cases} \tag{3.32}$$

$$|1 + R_o u_i + R_o u_i^2| = \begin{cases} p^{2r}, & \text{if char } R = p, \\ p^{3r}, & \text{if char } R = p^2, \\ p^{3r}, & \text{if char } R = p^3, \end{cases} \tag{3.33}$$

$$|1 + R_o u_i u_j| = p^r, \tag{3.34}$$

for every characteristic of  $R$ .

PROPOSITION 3.8. *The group  $1 + J$  is a direct product of the subgroup  $1 + pR_o$ ,  $s$  subgroups  $1 + R_o u_i + R_o u_i^2$ , and  $s(s - 1)/2$  subgroups  $1 + R_o u_i u_j$ , where  $i \neq j$  and  $u_i u_j = u_j u_i$ .*

*Proof.* This follows from the fact that  $1 + pR_o$ ,  $1 + R_o u_i + R_o u_i^2$ , and  $1 + R_o u_i u_j$  are subgroups of  $1 + J$ , intersection of any pair of these subgroups is trivial (for every  $i, j = 1, \dots, s$ ), and by Proposition 3.7,

$$|1 + J| = |1 + pR_o| \times \prod_{i=1}^s |1 + R_o u_i + R_o u_i^2| \times \prod_{i \neq j=1}^s |1 + R_o u_i u_j|. \tag{3.35}$$

□

3.2.1. *The structure of  $1 + pR_o$ .* The structure of  $1 + pR_o$  is completely determined by Raghavendran in [11]. For convenience of the reader, we state here the results useful for our purpose. For detailed proofs, refer to [11, Theorem 9].

We take  $r$  elements  $\varepsilon_1, \dots, \varepsilon_r$  in  $R_o$  with  $\varepsilon_1 = 1$  such that the set  $\{\overline{\varepsilon_1}, \dots, \overline{\varepsilon_r}\}$  is a basis of the quotient ring  $R_o/pR_o$  regarded as a vector space over its prime subfield  $GF(p)$ . Then we have the following.

PROPOSITION 3.9 [11, Theorem 9]. *If char  $R_o = p^2$ , then  $1 + pR_o$  is a direct product of  $r$  cyclic groups  $\langle 1 + p\varepsilon_j \rangle$ , each of order  $p$ , for any prime  $p$ .*

PROPOSITION 3.10 [11, Theorem 9]. *Let char  $R_o = p^3$ . If  $p = 2$ , then  $1 + pR_o$  is a direct product of 2 cyclic groups  $\langle -1 + 4\varepsilon_1 \rangle$  and  $\langle 1 + 4\varepsilon_1 \rangle$ , each of order 2, and  $(r - 1)$  cyclic groups  $\langle 1 + 2\varepsilon_j \rangle (j = 2, \dots, r)$ , each of order 4. If  $p \neq 2$ , then  $1 + pR_o$  is a direct product of  $r$  cyclic groups  $\langle 1 + p\varepsilon_j \rangle (j = 1, \dots, r)$ , each of order  $p^2$ .*

3.2.2. *The structure of  $1 + R_o u_i + R_o u_i^2$ .* We now consider the structure of the subgroup  $1 + R_o u_i + R_o u_i^2$  of the  $p$ -group  $1 + J$ . We first note that if char  $R_o = p$ , then  $R_o = GF(p^r)$  the field of  $p^r$  elements, if char  $R_o = p^2$ , then  $R_o$  is the Galois ring  $GR(p^{2r}, p^2)$  of order  $p^{2r}$ , and if char  $R_o = p^3$ ,  $R_o = GR(p^{3r}, p^3)$  the Galois ring of order  $p^{3r}$ .

We choose  $r$  elements  $\varepsilon_1, \dots, \varepsilon_r$  in  $R_o$  with  $\varepsilon_1 = 1$  such that the set  $\{\overline{\varepsilon_1}, \dots, \overline{\varepsilon_r}\}$  is a basis of the quotient ring  $R_o/pR_o$  regarded as a vector space over its prime subfield  $GF(p)$ . Then we have the following.

PROPOSITION 3.11. *Let  $\text{char } R_o = p$ . If  $p = 2$ , then  $1 + R_o u_i + R_o u_i^2$  is a direct product of  $r$  cyclic groups  $\langle 1 + \varepsilon_j u_i \rangle (j = 1, \dots, r)$ , each of order 4. If  $p \neq 2$ , then  $1 + R_o u_i + R_o u_i^2$  is a direct product of  $2r$  cyclic groups  $\langle 1 + \varepsilon_j u_i \rangle$  and  $\langle 1 + 2\varepsilon_j u_i \rangle (j = 1, \dots, r)$ , each of order  $p$ .*

*Proof.* If  $\text{char } R_o = 2$ , then  $\langle 1 + \varepsilon_j u_i \rangle$  is of order 4, for every  $j = 1, \dots, r$  and for any  $i = 1, \dots, s$ , and hence

$$\prod_{j=1}^r |\langle 1 + \varepsilon_j u_i \rangle| = 4^r = 2^{2r} = |1 + R_o u_i + R_o u_i^2|. \tag{3.36}$$

Therefore, the product  $\prod_{j=1}^r \langle 1 + \varepsilon_j u_i \rangle$  is direct.

Similarly, if  $\text{char } R_o = p \neq 2$ , the elements  $1 + \varepsilon_j u_i$  and  $1 + 2\varepsilon_j u_i$  are each of order  $p$ ,

$$\langle 1 + \varepsilon_j u_i \rangle \cap \langle 1 + 2\varepsilon_j u_i \rangle = \{1\}, \tag{3.37}$$

for every  $j = 1, \dots, r$ , and

$$\prod_{j=1}^r |\langle 1 + \varepsilon_j u_i \rangle| \cdot \prod_{j=1}^r |\langle 1 + 2\varepsilon_j u_i \rangle| = p^r \cdot p^r = p^{2r} = |1 + R_o u_i + R_o u_i^2|, \tag{3.38}$$

hence

$$1 + R_o u_i + R_o u_i^2 = \prod_{j=1}^r \langle 1 + \varepsilon_j u_i \rangle \times \prod_{j=1}^r \langle 1 + 2\varepsilon_j u_i \rangle, \tag{3.39}$$

a direct product. □

PROPOSITION 3.12. *Let  $\text{char } R_o = p^2$ . If  $p = 2$ , then  $1 + R_o u_i + R_o u_i^2$  is a direct product of  $r$  cyclic groups  $\langle 1 + 2\varepsilon_j u_i \rangle$ , each of order 2, and  $r$  cyclic groups  $\langle 1 + 3\varepsilon_j u_i \rangle (j = 1, \dots, r)$ , each of order 4. If  $p \neq 2$ , then  $1 + R_o u_i + R_o u_i^2$  is a direct product of  $r$  cyclic groups  $\langle 1 + p\varepsilon_j u_i \rangle$ , each of order  $p$ , and  $r$  cyclic groups  $\langle 1 + \varepsilon_j u_i \rangle (j = 1, \dots, r)$ , each of order  $p^2$ .*

*Proof.* Suppose  $\text{char } R_o = p^2$ . If  $p = 2$ ,  $\langle 1 + 2\varepsilon_j u_i \rangle$  is of order 2 and  $\langle 1 + 3\varepsilon_j u_i \rangle$  is of order 4,

$$\langle 1 + 2\varepsilon_j u_i \rangle \cap \langle 1 + 3\varepsilon_j u_i \rangle = \{1\}, \tag{3.40}$$

for every  $j = 1, \dots, r$  and any  $i = 1, \dots, s$ . Since

$$\prod_{j=1}^r |\langle 1 + 2\varepsilon_j u_i \rangle| \cdot \prod_{j=1}^r |\langle 1 + 3\varepsilon_j u_i \rangle| = 2^r \cdot 4^r = 2^{3r} = |1 + R_o u_i + R_o u_i^2|, \tag{3.41}$$

it follows that

$$1 + R_o u_i + R_o u_i^2 = \prod_{j=1}^r \langle 1 + 2\varepsilon_j u_i \rangle \times \prod_{j=1}^r \langle 1 + 3\varepsilon_j u_i \rangle \tag{3.42}$$

is a direct product.

If  $p \neq 2$ , it is easy to check that  $|\langle 1 + p\varepsilon_j u_i \rangle| = p$ ,  $|\langle 1 + \varepsilon_j u_i \rangle| = p^2$  and

$$\langle 1 + p\varepsilon_j u_i \rangle \cap \langle 1 + \varepsilon_j u_i \rangle = \{1\}, \tag{3.43}$$

for every  $j = 1, \dots, r$  and any  $i = 1, \dots, s$ . Since

$$\prod_{j=1}^r |\langle 1 + p\varepsilon_j u_i \rangle| \cdot \prod_{j=1}^r |\langle 1 + \varepsilon_j u_i \rangle| = p^r \cdot (p^2)^r = p^{3r} = |1 + R_o u_i + R_o u_i^2|, \tag{3.44}$$

it follows that the product

$$1 + R_o u_i + R_o u_i^2 = \prod_{j=1}^r \langle 1 + 2\varepsilon_j u_i \rangle \times \prod_{j=1}^r \langle 1 + 3\varepsilon_j u_i \rangle \tag{3.45}$$

is direct. □

**PROPOSITION 3.13.** *Let  $\text{char } R_o = p^3$ . If  $p = 2$ , then  $1 + R_o u_i + R_o u_i^2$  is a direct product of  $r$  cyclic groups  $\langle 1 + \varepsilon_j u_i^2 \rangle$ , each of order 2, and  $r$  cyclic groups  $\langle 1 + \varepsilon_j u_i \rangle$  ( $j = 1, \dots, r$ ), each of order 4. If  $p \neq 2$ , then  $1 + R_o u_i + R_o u_i^2$  is a direct product of  $r$  cyclic groups  $\langle 1 + \varepsilon_j u_i^2 \rangle$ , each of order  $p$ , and  $r$  cyclic groups  $\langle 1 + \varepsilon_j u_i \rangle$  ( $j = 1, \dots, r$ ), each of order  $p^2$ .*

*Proof.* Similar to the proofs of Propositions 3.11 and 3.12. □

**3.2.3. The structure of  $1 + R_o u_i u_j$ .** Choose  $r$  elements  $\varepsilon_1, \dots, \varepsilon_r$  in  $R_o$  with  $\varepsilon_1 = 1$  such that the elements  $\bar{\varepsilon}_1, \dots, \bar{\varepsilon}_r$  form a basis of the quotient ring  $R_o/pR_o$  regarded as a vector space over its prime subfield  $GF(p)$ . Then we have the following.

**PROPOSITION 3.14.** *The group  $1 + R_o u_i u_j$  is a direct product of  $r$  cyclic groups  $\langle 1 + \varepsilon_l u_i u_j \rangle$  ( $l = 1, \dots, r$ ), each of order  $p$ , for any characteristic  $p^k$  ( $1 \leq k \leq 3$ ) of  $R$ .*

*Proof.* We first note that if the characteristic of  $R$  is  $p^k$ , where  $1 \leq k \leq 3$ , then  $pu_i u_j = 0$ . Hence,  $|1 + R_o u_i u_j| = p^r$ . Also, for any  $x \in 1 + R_o u_i u_j$ ,  $x^p = 1$ .

Now, for  $r$  elements  $\varepsilon_1, \dots, \varepsilon_r \in R_o$  defined above, since for any  $\nu \neq \mu$ ,

$$\langle 1 + \varepsilon_\nu u_i u_j \rangle \cap \langle 1 + \varepsilon_\mu u_i u_j \rangle = 1, \tag{3.46}$$

the result follows. □

We now state the main results of this section.

**THEOREM 3.15.** *Let  $\text{char } R = p$ . If  $p = 2$ , then  $1 + J$  is a direct product of  $(s(s - 1)/2)r$  cyclic groups, each of order 2, and  $sr$  cyclic groups, each of order 4. If  $p \neq 2$ , then  $1 + J$  is a direct product of  $((s^2 + 3s)/2)r$  cyclic groups, each of order  $p$ .*

*Proof.* This follows from Propositions 3.11 and 3.14 and by the fact that the order of  $1 + J$  is  $p^{((s^2+3s)/2)r}$ . □

**THEOREM 3.16.** *Let  $\text{char } R = p^2$ . Then  $1 + J$  is a direct product of  $((s^2 + s + 2)/2)r$  cyclic groups, each of order  $p$ , and  $sr$  cyclic groups, each of order  $p^2$ , for any prime  $p$ .*

*Proof.* This follows from Propositions 3.9, 3.12, and 3.14 and from the fact that the order of  $1 + J$  is  $p^{((s^2+5s+2)/2)r}$ . □

**THEOREM 3.17.** *Let  $\text{char } R = p^3$ . If  $p = 2$ , then  $1 + J$  is a direct product of  $2 + ((s^2 + s)/2)r$  cyclic groups, each of order 2, and  $r - 1 + sr$  cyclic groups, each of order 4. If  $p \neq 2$ , then  $1 + J$  is a direct product of  $((s^2 + s)/2)r$  cyclic groups, each of order  $p$ , and  $(s + 1)r$  cyclic groups, each of order  $p^2$ .*

*Proof.* First observe that the order of  $1 + J$  is  $p^{((s^2+5s+4)/2)r}$ . By Propositions 3.10, 3.13, and 3.14, the result follows. □

**4. The Main theorem**

By Proposition 2.1, the group of units  $R^*$  of  $R$  contains a cyclic subgroup  $\langle b \rangle$  of order  $p^r - 1$ , and  $R^*$  is a direct product of  $1 + J$  and  $\langle b \rangle$ . Moreover, the structure of  $1 + J$  has been determined in Section 3 (Theorems 3.15, 3.16, and 3.17). We thus have the following result.

**THEOREM 4.1.** *The group of units  $R^*$ , of a commutative completely primary finite ring  $R$  with maximal ideal  $J$  such that  $J^3 = (0)$  and  $J^2 \neq (0)$ , and with invariants  $p, k, r, s$ , and  $t$ , where  $t = s(s + 1)/2$ , is a direct product of cyclic groups as follows:*

(i) if  $\text{char } R = p$ , then

$$R^* \cong \begin{cases} \mathbb{Z}_{2^{r-1}} \times (\mathbb{Z}_4^r)^s \times (\mathbb{Z}_2^r)^\gamma, & \text{if } p = 2, \\ \mathbb{Z}_{p^{r-1}} \times (\mathbb{Z}_p^r)^s \times (\mathbb{Z}_p^r)^\gamma, & \text{if } p \neq 2, \end{cases} \tag{4.1}$$

(ii) if  $\text{char } R = p^2$ , then

$$R^* \cong \begin{cases} \mathbb{Z}_{2^{r-1}} \times \mathbb{Z}_2^r \times (\mathbb{Z}_2^r)^s \times (\mathbb{Z}_2^r)^\gamma, & \text{if } p = 2, \\ \mathbb{Z}_{p^{r-1}} \times \mathbb{Z}_p^r \times (\mathbb{Z}_p^r)^s \times (\mathbb{Z}_{p^2}^r)^\gamma, & \text{if } p \neq 2, \end{cases} \tag{4.2}$$

(iii) if  $\text{char } R = p^3$ , then

$$R^* \cong \begin{cases} \mathbb{Z}_{2^{r-1}} \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4^{r-1} \times (\mathbb{Z}_2^r)^s \times (\mathbb{Z}_4^r)^s \times (\mathbb{Z}_2^r)^\gamma, & \text{if } p = 2, \\ \mathbb{Z}_{p^{r-1}} \times \mathbb{Z}_{p^2}^r \times (\mathbb{Z}_p^r)^s \times (\mathbb{Z}_{p^2}^r)^\gamma, & \text{if } p \neq 2, \end{cases} \tag{4.3}$$

where  $\gamma = (s^2 - s)/2$ .

*Proof.* Follows from Propositions 2.1 and 3.9 through 3.14 and Theorems 3.15, 3.16, and 3.17. □




*Remark 4.2.* The structure of the multiplicative groups of commutative completely primary finite rings  $R$  with maximal ideals  $J$  such that  $J^3 = (0)$  and  $J^2 \neq (0)$ , for which  $t < s(s + 1)/2$  for a fixed nonnegative integer  $s$ , will be considered in subsequent work.

**References**

- [1] C. W. Ayoub, *On finite primary rings and their groups of units*, *Compositio Math.* **21** (1969), 247–252.
- [2] C. J. Chikunji, *On a class of finite rings*, *Comm. Algebra* **27** (1999), no. 10, 5049–5081.
- [3] ———, *On a class of rings of order  $p^5$* , *Math. J. Okayama Univ.* **45** (2003), 59–71.
- [4] W. E. Clark, *A coefficient ring for finite non-commutative rings*, *Proc. Amer. Math. Soc.* **33** (1972), 25–28.
- [5] D. Dolzan, *Group of units in a finite ring*, *J. Pure Appl. Algebra* **170** (2002), no. 2-3, 175–183.
- [6] L. Fuchs, *Abelian Groups*, 3rd ed., *International Series of Monographs on Pure and Applied Mathematics*, Pergamon Press, New York, 1960.
- [7] G. Ganske and B. R. McDonald, *Finite local rings*, *Rocky Mountain J. Math.* **3** (1973), no. 4, 521–540.
- [8] R. W. Gilmer Jr., *Finite rings having a cyclic multiplicative group of units*, *Amer. J. Math.* **85** (1963), 447–452.
- [9] D. Gorenstein, R. Lyons, and R. Solomon, *The Classification of the Finite Simple Groups*, vol. 40, *Mathematical Surveys and Monographs*, no. 1, American Mathematical Society, Rhode Island, 1994.
- [10] K. R. Pearson and J. E. Schneider, *Rings with a cyclic group of units*, *J. Algebra* **16** (1970), 243–251.
- [11] R. Raghavendran, *Finite associative rings*, *Compositio Math.* **21** (1969), 195–229.
- [12] I. Stewart, *Finite rings with a specified group of units*, *Math. Z.* **126** (1972), 51–58.

Chiteng'a John Chikunji: Department of Mathematics, University of Transkei, Private Bag X1, Umtata 5117, South Africa

*E-mail address:* chikunji@getafix.ut.ac.za



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

